

Two doses a day are preferred to one in order to maintain as smooth a diurnal tension curve as possible, although a single dose per day or every other day has been used with satisfactory results. The daily dose or one of the two daily doses should always be instilled just before retiring to avoid inconvenience due to the miosis.

Technique – Technique in the administration of echothiophate iodide for ophthalmic solution may include finger pressure at the inner canthus exerted for a minute or two following instillation of the eyedrops, to potentially minimize drainage into the nose and throat. Excess solution around the eye should be removed with tissue and any medication on the hands should be rinsed off.

Accommodative Esotropia (Pediatric Use)

In Diagnosis – One drop may be instilled once a day in both eyes on retiring, for a period of two or three weeks. If the esotropia is accommodative, a favorable response will usually be noted which may begin within a few hours.

In Treatment – Echothiophate iodide for ophthalmic solution is prescribed at the lowest frequency which gives satisfactory results. After the initial period of treatment for diagnostic purposes, the schedule may be reduced to one drop every other day. The maximum usually recommended dosage is one drop once a day, although more intensive therapy has been used for short periods.

Technique – (See “**DOSAGE AND ADMINISTRATION**”)

Duration of Treatment — In therapy, there is no definite limit so long as the drug is well tolerated. However, if the eyedrops, with or without eyeglasses, are gradually withdrawn after about a year or two and deviation recurs, surgery should be considered. As with other miotics, tolerance may occasionally develop after prolonged use. In such cases, a rest period may restore the original activity of the drug.

HOW SUPPLIED

Each package contains sterile echothiophate iodide for ophthalmic solution, sterile diluent, and dropper for dispensing 5 mL eyedrops.

NDC 48102-053-05 6.25 mg package for 0.125%
White amorphous deposit on bottle walls. Aluminum crimp seal is green.

HANDLING AND STORAGE:

Prior to reconstitution: Store under refrigeration (2°C to 8°C).

After reconstitution: Store upright at room temperature (approximately 25°C) (68°F to 77°F). Do not refrigerate. Discard any unused solution after 4 weeks.



Distributed by:
Fera Pharmaceuticals, LLC
Locust Valley, NY 11560

PF053
Rev. 0224

ART: 3 DATE: 02/15/2024 CUSTOMER: FERA			<div>APPROVED</div> <div>(Signature / Date)</div>	NON PRINTING	SPOT INKS	SPECIAL FEATURES	<div>ArtFlo</div> <div>by Nosco®</div>			
P/N: Phospholine-Iodide_PF053_0224_PI				PROCESS INKS						
Eval#: X5823324			<div>Black</div>			COATING				
SIZE: 15.0 x 6.625 FOLD: 3.0 x 1.125 DRWG: 231548										
<table><tr><td>Barcode Type</td><td>Decoded Data</td><td>Human Readable</td></tr></table>			Barcode Type	Decoded Data	Human Readable					
Barcode Type	Decoded Data	Human Readable								
Codes:										
<div>• CAUTION Artwork cycles are NOT proofread. Please, proofread BEFORE authorizing Nosco to proceed to final proofing. This ART is to show size, copy placement and color breaks. Actual colors will be matched on press to approved color standards and/or PMS color swatches. *Customer Logo's, Brand/Drug names with SuperscriptsSubscripts with ®s and TM's is to be a minimum 1/32 high for the TM and bold, and for the® 1/32 in circumference and bold. Any deviation to this will be printed as is. Nosco Glue Flap Barcode: is for internal Quality Control Purposes Only and any element in Die color Does NOT Print.</div>										